

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/043006

International filing date: 17 December 2004 (17.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/534,442
Filing date: 06 January 2004 (06.01.2004)

Date of receipt at the International Bureau: 17 January 2005 (17.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

January 10, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/534,442

FILING DATE: *January 06, 2004*

RELATED PCT APPLICATION NUMBER: *PCT/US04/43006*



Certified By

Jon W Dudas

Under Secretary
of Commerce for Intellectual Property
and Acting Director of the
United States Patent and Trademark Office

Please type a plus sign (+) inside this box

+

PTO/SB/16 (5-03)
Approved for use through 04/30/2003. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
BINH THANH		NGUYEN		MIDLAND, MICHIGAN 48640 USA	
<input type="checkbox"/> Additional inventors are being named on the _____ separately numbered sheets attached hereto					
TITLE OF THE INVENTION (280 characters max)					
GRIGNARD PROCESSES WITH INCREASED CONTENT OF DIPHENYLCHLOROSILANES					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input checked="" type="checkbox"/> Customer Number		00137		<input type="checkbox"/> Place Customer Number Bar Code Label here	
OR Type Customer Number here					
<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		ZIP	
Country		Telephone		Fax	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification		Number of Pages		9	
<input type="checkbox"/> Drawing(s)		Number of Sheets			
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76		<input type="checkbox"/> CD(s), Number			
		<input checked="" type="checkbox"/> Other (specify)		3 PAGES CLAIMS 1 PAGE ABSTRACT	
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)					
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees				FILING FEE AMOUNT (\$)	
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number		04-1520		\$160.00	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.					
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

Respectfully submitted,

SIGNATURE



TYPED or PRINTED NAME JIM DECESARE

TELEPHONE

(989) 496-4235

Date

01/05/2004

REGISTRATION NO.

27,979


(if appropriate)

Docket Number:

DC10001PSP1

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

CERTIFICATE OF MAILING BY "EXPRESS MAIL" (37 CFR 1.10) Applicant(s): BINH THANH NGUYEN			Docket No. DC10001PSP1
Serial No. TO BE ASSIGNED	Filing Date TO BE ASSIGNED	Examiner TO BE ASSIGNED	Group Art Unit TO BE ASSIGNED
Invention: GRIGNARD PROCESSES WITH INCREASED CONTENT OF DIPHENYLCHLOROSILANES			
<p>I hereby certify that the following correspondence:</p> <div style="border: 1px solid black; padding: 5px; text-align: center;">PROVISIONAL APPLICATION FOR PATENT COVER SHEET; APPLICATION; POSTCARD</div> <p style="text-align: center;"><i>(Identify type of correspondence)</i></p> <p>is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450 on</p> <p style="text-align: center;"><u>January 6, 2004</u> <i>(Date)</i></p> <div style="text-align: center;"><p>KELLY J. SMITH <i>(Typed or Printed Name of Person Mailing Correspondence)</i></p><p> <i>(Signature of Person Mailing Correspondence)</i></p><p>EV319568587US <i>("Express Mail" Mailing Label Number)</i></p></div>			
<p>Note: Each paper must have its own certificate of mailing.</p>			

UNITED STATES PATENT APPLICATION

TITLE

Grignard Processes With Increased Content of Diphenylchlorosilanes

5

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] Not applicable.

FIELD OF THE INVENTION

- 10 [0002] This invention is related to Grignard processes for preparing phenyl-containing chlorosilane products. In particular, it relates to Grignard processes in which the yield of diphenylchlorosilanes as a product is maximized, and the yield of phenylchlorosilanes as a product is minimized.

15 BACKGROUND OF THE INVENTION

- [0003] In copending US Patent Application Serial No. 10/117,259, filed on April 4, 2002, entitled "Process for Preparing Phenylorganosilicon Intermediates" (the '259 application hereafter), assigned to the same assignee as the present invention, a Grignard process is described in which the yield of phenylchlorosilanes as a product is maximized and the yield of diphenylchlorosilanes as a product is minimized. In fact, the diphenylchlorosilanes prepared in the '651 patent are only present as by-products.

20

- [0004] This invention in contrast, is characterized in that it seeks to obtain an opposite result, i.e., to minimize the yield of phenylchlorosilanes as a product while maximizing the yield of diphenylchlorosilanes as a product. Achievement of this goal is obtained by carrying out the Grignard process using certain mole ratios of the reactants used in the Grignard process.

25

SUMMARY OF THE INVENTION

- [0005] This invention is directed to three improved Grignard processes for preparing phenyl-containing chlorosilane products in which the yield of diphenylchlorosilanes as a product is maximized, and the yield of phenylchlorosilanes as a product is minimized.

30

[0006] In a first normal coupling process embodiment, the process involves contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, and a trichlorosilane, in a mole ratio of 1/4/3/0.5, respectively.

[0007] In a second co-coupling process embodiment, the process involves contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, a trichlorosilane, and a phenylchlorosilane, in a mole ratio of 1/4/3/1.3/0.38, respectively.

[0008] In a third direct coupling process embodiment, the process involves contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, and a phenylchlorosilane, in a mole ratio of 1/4/3/1.1, respectively.

[0009] In the first embodiment, the mole ratio of the ether solvent to the phenyl Grignard reagent is 2 to 5, the mole ratio of the aromatic halogenated coupling solvent to the phenyl Grignard reagent is 3 to 7, and the mole ratio of the trichlorosilane to the phenyl Grignard reagent is 0.1 to 10. In the second embodiment, the mole ratio of the ether solvent to the phenyl Grignard reagent is 2 to 5, the mole ratio of the aromatic halogenated coupling solvent to the phenyl Grignard reagent is 3 to 7, the mole ratio of the trichlorosilane to the phenyl Grignard reagent is 0.1 to 10, and the mole ratio of the phenylchlorosilane to the phenyl Grignard reagent is 0.5 to 5. In the third embodiment, the mole ratio of the ether solvent to the phenyl Grignard reagent is 2 to 5, the mole ratio of the aromatic halogenated coupling solvent to the phenyl Grignard reagent is 3 to 7, and the mole ratio of the phenylchlorosilane to the phenyl Grignard reagent is 0.5 to 5.

[0010] These and other features of the invention will become apparent from a consideration of the detailed description.



BRIEF DESCRIPTION OF THE DRAWING

[0011] Not applicable.

DETAILED DESCRIPTION OF THE INVENTION



[0012] As used herein, the term *normal coupling* refers to reactions of a phenyl Grignard reagent chloride with a trichlorosilane; the term *co-coupling* refers to reactions of the phenyl Grignard reagent the trichlorosilane and a phenylchlorosilane; and the term *direct coupling* refers to reactions of the phenyl Grignard reagent with the phenylchlorosilane

[0013] The Grignard process according to this invention is illustrated below in chemical reactions (I) and (II). Chemical reaction (II) depicts the first embodiment of the invention (normal coupling). Chlorobenzene is also one of the products in chemical reaction (II) but is not shown in chemical reaction (II).




[0014] In chemical reaction (I), phenyl chloride/chlorobenzene (PhCl) is combined with magnesium metal (Mg) in the presence of the solvent diethyl ether ($\text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_3$), to form phenyl magnesium chloride (PhMgCl) in diethyl ether. Phenyl magnesium chloride in diethyl ether is then used in chemical reaction (II) where it is combined with methyltrichlorosilane (MeSiCl_3) and the coupling solvent chlorobenzene. The products of chemical reaction (II) are phenylmethyldichlorosilane (PhMeSiCl_2), diphenylmethylchlorosilane (Ph_2MeSiCl), magnesium chloride, and chlorobenzene.

[0015] According to the second embodiment of the invention (co-coupling) depicted below as chemical reaction (III), phenylmethyldichlorosilane is added as an additional reactant along with methyltrichlorosilane. Chlorobenzene is also one of the products in chemical reaction (III) but is not shown in chemical reaction (III).



[0016] The third embodiment of the invention (direct coupling) is similar to the first embodiment of chemical reaction (II), except that in the third embodiment in chemical reaction (IV), phenylmethyldichlorosilane is used instead of methyltrichlorosilane.

Chlorobenzene is also one of the products in chemical reaction (IV) but is not shown in chemical reaction (IV).



[0017] In the first embodiment of this invention (normal coupling), the mole ratio of PhMgCl /ether/chlorobenzene/ MeSiCl_3 is 1/4/3/0.5, respectively, as compared to the corresponding mole ratio of PhMgCl /ether/chlorobenzene/ MeSiCl_3 in the '259 application of 1/0.08/4/2, in Example 3.

[0018] In the second embodiment of this invention (co-coupling), the mole ratio of PhMgCl /ether/chlorobenzene/ MeSiCl_3 / PhMeSiCl_2 is 1/4/3/1.3/0.38, respectively. This co-coupling method, however, is not even disclosed in the '259 application.

[0019] In the third embodiment of this invention (direct coupling), the mole ratio of PhMgCl/ether/chlorobenzene//PhMeSiCl₂ is 1/4/3/1.1, respectively. This direct coupling method, however, is not even disclosed in the '259 application.

[0020] It should be noted that there are significant differences between the three processes described in this invention in comparison to the single process described in the common assignee's '259 application. Thus, the mole ratio according to the first embodiment of this invention (normal coupling) is different than the mole ratio in the '259 application, and this results in an almost three-fold increase in the yield of diphenylmethylchlorosilane in comparison to the yield of only about 6 percent obtained in the '259 application. In addition, neither of the processes of the second embodiment (co-coupling) and of the third embodiments (direct coupling) are even present in the '259 application.

[0021] These differences are also significant in that the thrust and focus of the '259 application is the production of phenylmethyldichlorosilane as the primary and major product of the reaction, whereas the thrust and focus of the present invention is the production of diphenylmethylchlorosilane as the desirable product of the three processes. Thus, by following the teaching of the method according to the first embodiment of this invention, it is possible for one skilled in the art to prepare products of the process containing about 20 percent or more by weight of diphenylmethylchlorosilane, as compared to the products obtained in the '259 application which contain only about 1-6 percent by weight of phenylmethyldichlorosilane. The '259 application is completely silent regarding the second and third process embodiments of this invention.

[0022] Chlorosilanes useful according to the invention have the general formula R_aSiX_{4-a} wherein each R can represent a phenyl group, methyl group, vinyl group, or hydrogen; X represents chlorine or bromine; and a has a value of 0, 1, or 2. Some suitable and representative chlorosilanes which can be used include silicon tetrachloride, methyltrichlorosilane, dimethyldichlorosilane, phenylmethyldichlorosilane, phenyltrichlorosilane, diphenyldichlorosilane, vinyltrichlorosilane, hydridotrichlorosilane, divinylchlorosilane, methylvinylchlorosilane, phenylvinylchlorosilane, hydridomethylchlorosilane, hydridophenylchlorosilane, hydridovinylchlorosilane and dihydridodichlorosilane.

[0023] Magnesium metal useful in this invention can be any of the forms of the metal currently being used in Grignard-type reactions. For example, the metal can be in the form of

a powder, flake, granule, chip, lump, or shaving. Contact of the magnesium metal with the phenyl halide can be undertaken in standard type reactors suitable for running Grignard type reactions. Thus, the reactor can be a batch, semi-batch, or continuous type reactor. A preferred reactor is a continuous reactor. The environment in which the present method is carried out should be inert for best results. Therefore, under preferred conditions of the method, the reactor is purged and blanketed with an inert gas such as nitrogen or argon.

[0024] Phenyl halides useful in this invention are those of the formula RX wherein R represents phenyl and X is a chlorine or bromine atom. The preferred phenyl halide for this invention is phenyl chloride (chlorobenzene). Solvents for synthesizing the Grignard reagent include dialkyl ethers such as dimethyl ether, diethyl ether, ethylmethyl ether, n-butylmethyl ether, n-butylethyl ether, di-n-butyl ether, di-isobutyl ether, isobutylmethyl ether, and isobutylethyl ether. The most preferred ether solvent is diethyl ether. Aromatic halogenated solvents such as chlorobenzene and 1,4-dichlorobenzene are used as the coupling solvent in the coupling reaction of the phenyl Grignard reagent $PhMgCl$ with $MeSiCl_3$, mixtures of $MeSiCl_3$ and $PhMeSiCl_2$, or $PhMeSiCl_2$, according to the processes of the invention. Phenyl Grignard reagents such as $PhMgCl$ can either be synthesized or purchased commercially, as desired.

EXAMPLES

[0025] The following examples are set forth in order to illustrate the invention in more detail.

Example 1 - Normal Coupling Method

[0026] In this example, the coupling mole ratio was 1/4/3/0.5 of phenylmagnesium chloride/diethyl ether/chlorobenzene/methyltrichlorosilane ($PhMgCl/Et_2O/C_6H_5Cl/MeSiCl_3$). The $MeSiCl_3$ and the Grignard solution ($PhMgCl/Et_2O$) used in this example were compositions prepared by an in-house facility. The mole ratio of the Grignard solution was 1/4 $PhMgCl/Et_2O$ with a $PhMgCl$ concentration of 2 mol/L. This Grignard solution contained two phases, a liquid phase and a solids phase that settled at the bottom. Both the liquid phase and the solid phase were used in this example. Approximately 250 milliliter of the Grignard solution were transferred to a 500 milliliter addition funnel by means of a pump. The amount of Grignard solution used was the equivalent of about 0.500

mol of PhMgCl and 2.000 mol of Et₂O. Then, 37.37 gram (0.25 mol) of MeSiCl₃ and 168.75 gram (1.499 mol) of PhCl were added to a 1000 milliliter round bottom flask.

[0027] The addition funnel was connected to a source of nitrogen to provide an inert atmosphere for the system and placed on a 500 milliliter three-neck round bottom flask containing a magnetic stirrer. The 500 milliliter three-neck round bottom was charged with the Grignard solution and stirred. From the 500 milliliter three-neck round bottom flask, the Grignard solution was pumped into a reactor. Addition of the Grignard solution took place over a time period of about ten minutes. The solution turned dark orange brown in color but remained in a flowable state throughout the procedure. The maximum exothermic temperature reached was 64 °C. When the agitation was stopped, the settling of solids began almost immediately. The reaction mixture was allowed to cool and then it was transferred to a labeled sample jar. The percent mass recovery was determined to be about 94.90 percent.

[0028] Gas Chromatography (GC) analysis of the reaction mixture showed that it contained about 20.8 weight percent of the desired component Ph₂MeSiCl, and only a small amount, i.e., 0.07 weight percent, of PhMeSiCl₂ as a by-product.

Example 2 - Co-Coupling Method

[0029] In this example, the coupling mole ratio was 1/4/3/1.3/0.38 of phenylmagnesium chloride/diethyl ether/chlorobenzene/methyltrichlorosilane/phenylmethyldichlorosilane (PhMgCl/Et₂O/C₆H₅Cl/MeSiCl₃/PhMeSiCl₂). The MeSiCl₃ and the Grignard solution (PhMgCl/Et₂O) used in this example were compositions prepared by an in-house facility. The mole ratio of the Grignard solution was 1/4 PhMgCl/Et₂O with a PhMgCl concentration of 2 mol/L. This Grignard solution contained two phases, a liquid phase and a solids phase that settled at the bottom. Both the liquid phase and the solid phase were used in this example. Approximately 250 milliliter of the Grignard solution were transferred to a 500 milliliter addition funnel by means of a pump. The amount of Grignard solution used was the equivalent of about 0.500 mol of PhMgCl and 2.000 mol of Et₂O. Then, 99.41 gram (0.67 mol) of MeSiCl₃, 36.32 gram (0.19 mol) of PhMeSiCl₂, and 168.75 gram (1.499 mol) of PhCl were added to a 1000 milliliter round bottom flask.

[0030] The addition funnel was connected to a source of nitrogen to provide an inert atmosphere for the system and placed on a 500 milliliter three-neck round bottom flask

containing a magnetic stirrer. The 500 milliliter three-neck round bottom was charged with the Grignard solution and stirred. From the 500 milliliter three-neck round bottom flask, the Grignard solution was pumped into a reactor. Addition of the Grignard solution took place over a time period of about ten minutes. The solution turned dark orange brown in color but remained in a flowable state throughout the procedure. The maximum exothermic temperature reached was 67 °C. When the agitation was stopped, the settling of solids began almost immediately. The reaction mixture was allowed to cool and then it was transferred to a labeled sample jar. The percent mass recovery was determined to be about 94.11 percent.

[0031] Gas Chromatography (GC) analysis of the reaction mixture showed that it contained about 2.9 weight percent of the desired component Ph_2MeSiCl , and about 29.7 weight percent of PhMeSiCl_2 . In this co-coupling method, the weight ratio of $\text{PhMeSiCl}_2/\text{Ph}_2\text{MeSiCl}$ was 0.1.

Example 3 - Direct Coupling Method

[0032] In this example, the coupling mole ratio was 1/4/3/1.1 of phenylmagnesium chloride/diethyl ether/chlorobenzene/phenylmethyldichlorosilane ($\text{PhMgCl}/\text{Et}_2\text{O}/\text{C}_6\text{H}_5\text{Cl}/\text{PhMeSiCl}_2$). The MeSiCl_3 and the Grignard solution ($\text{PhMgCl}/\text{Et}_2\text{O}$) used in this example were compositions prepared by an in-house facility. The mole ratio of the Grignard solution was 1/4 $\text{PhMgCl}/\text{Et}_2\text{O}$ with a PhMgCl concentration of 2 mol/L. This Grignard solution contained two phases, a liquid phase and a solids phase that settled at the bottom. Both the liquid phase and the solid phase were used in this example. Approximately 250 milliliter of the Grignard solution were transferred to a 500 milliliter addition funnel by means of a pump. The amount of Grignard solution used was the equivalent of about 0.500 mol of PhMgCl and 2.000 mol of Et_2O . Then, 105.127 gram (0.55 mol) of PhMeSiCl_2 and 168.75 gram (1.499 mol) of PhCl were added to a 1000 milliliter round bottom flask.

[0033] The addition funnel was connected to a source of nitrogen to provide an inert atmosphere for the system and placed on a 500 milliliter three-neck round bottom flask containing a magnetic stirrer. The 500 milliliter three-neck round bottom was charged with the Grignard solution and stirred. From the 500 milliliter three-neck round bottom flask, the Grignard solution was pumped into a reactor. Addition of the Grignard solution took place

over a time period of about 10.5 minutes. The solution turned dark orange brown in color but remained in a flowable state throughout the procedure. The maximum exothermic temperature reached was 62 °C. When the agitation was stopped, the settling of solids began almost immediately. The reaction mixture was allowed to cool and then it was transferred to a labeled sample jar. The percent mass recovery was determined to be about 92.11 percent.

[0034] Gas Chromatography (GC) analysis of the reaction mixture showed that it contained about 12.18 weight percent of the desired component Ph_2MeSiCl , and about 3.6 weight percent of PhMeSiCl_2 . In this direct coupling method, the weight ratio of $\text{PhMeSiCl}_2/\text{Ph}_2\text{MeSiCl}$ was 3.4.

[0035] Those skilled in the art are cognizant of the fact that it is both difficult and not easily obtainable, to increase the formation of Ph_2MeSiCl by a normal coupling reaction of PhMgCl with MeSiCl_3 , or by a co-coupling reaction of PhMgCl with MeSiCl_3 and PhMeSiCl_2 , or by a direct coupling reaction of PhMgCl with PhMeSiCl_2 . Contrary to the art however, it has been unexpectedly discovered according to this invention, that by changing the ratio of $\text{PhMgCl}/\text{MeSiCl}_3$ (normal-coupling), or $\text{PhMgCl}/\text{MeSiCl}_3/\text{PhMeSiCl}_2$ (co-coupling), or $\text{PhMgCl}/\text{PhMeSiCl}_2$ (direct coupling), it is possible to not only improve, but to actually increase the production of Ph_2MeSiCl . This is significant, since it now enables those skilled in the art to increase and maximize production of Ph_2MeSiCl , while at the same time minimizing production of PhMeSiCl_2 , by a process not heretofore known.

[0036] Other variations may be made in compounds, compositions, and methods described herein without departing from the essential features of the invention. The embodiments of the invention specifically illustrated herein are exemplary only and not intended as limitations on their scope except as defined in the appended claims.

CLAIM OR CLAIMS

1. A process for preparing diphenylchlorosilanes by the Grignard process comprising contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, and a trichlorosilane; wherein the mole ratio of the ether solvent to the phenyl Grignard reagent is 2 to 5, the mole ratio of the aromatic halogenated coupling solvent to the phenyl Grignard reagent is 3 to 7, and the mole ratio of the trichlorosilane to the phenyl Grignard reagent is 0.1 to 10.
2. The process according to Claim 1 wherein the phenyl Grignard reagent is phenyl magnesium chloride.
3. The process according to Claim 1 wherein the ether solvent is a dialkyl ether selected from the group consisting of dimethyl ether, diethyl ether, ethylmethyl ether, n-butylmethyl ether, n-butylethyl ether, di-n-butyl ether, di-isobutyl ether, isobutylmethyl ether, and isobutylethyl ether.
4. The process according to Claim 1 wherein the aromatic halogenated coupling solvent is chlorobenzene.
5. The process according to Claim 1 wherein the trichlorosilane is selected from the group consisting of methyltrichlorosilane, phenyltrichlorosilane, and vinyltrichlorosilane.
6. A process for preparing diphenylchlorosilanes by the Grignard process comprising contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, a trichlorosilane, and a phenylchlorosilane; wherein the mole ratio of the ether solvent to the phenyl Grignard reagent is 2 to 5, the mole ratio of the aromatic halogenated coupling solvent to the phenyl Grignard reagent is 3 to 7, the mole ratio of the trichlorosilane to the phenyl Grignard reagent is 0.1 to 10, and the mole ratio of the phenylchlorosilane to the phenyl Grignard reagent is 0.5 to 5.

7. The process according to Claim 6 wherein the phenyl Grignard reagent is phenyl magnesium chloride.
8. The process according to Claim 6 wherein the ether solvent is a dialkyl ether selected from the group consisting of dimethyl ether, diethyl ether, ethylmethyl ether, n-butylmethyl ether, n-butyloethyl ether, di-n-butyl ether, di-isobutyl ether, isobutylmethyl ether, and isobutyloethyl ether.
9. The process according to Claim 6 wherein the aromatic halogenated coupling solvent is chlorobenzene.
10. The process according to Claim 6 wherein the trichlorosilane is selected from the group consisting of methyltrichlorosilane, phenyltrichlorosilane, and vinyltrichlorosilane.
11. The process according to Claim 6 wherein the phenylchlorosilane is selected from the group consisting of phenylmethyldichlorosilane, phenyltrichlorosilane, diphenyldichlorosilane, phenylvinylchlorosilane, and hydridophenyldichlorosilane.
12. A process for preparing diphenylchlorosilanes by the Grignard process comprising contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, and a phenylchlorosilane; wherein the mole ratio of the ether solvent to the phenyl Grignard reagent is 2 to 5, the mole ratio of the aromatic halogenated coupling solvent to the phenyl Grignard reagent is 3 to 7, and the mole ratio of the phenylchlorosilane to the phenyl Grignard reagent is 0.5 to 5,
13. The process according to Claim 12 wherein the phenyl Grignard reagent is phenyl magnesium chloride.
14. The process according to Claim 12 wherein the ether solvent is a dialkyl ether selected from the group consisting of dimethyl ether, diethyl ether, ethylmethyl ether, n-butylmethyl ether, n-butyloethyl ether, di-n-butyl ether, di-isobutyl ether, isobutylmethyl ether, and isobutyloethyl ether.

15. The process according to Claim 12 wherein the aromatic halogenated coupling solvent is chlorobenzene.

- 5 16. The process according to Claim 12 wherein the phenylchlorosilane is selected from the group consisting of phenylmethyldichlorosilane, phenyltrichlorosilane, diphenyldichlorosilane, phenylvinylidichlorosilane, and hydridophenyldichlorosilane.

10

15

20

25

30

ABSTRACT

Three improved Grignard processes are used for preparing phenyl-containing chlorosilane products wherein the yield of diphenylchlorosilanes as a product is maximized, while the yield of phenylchlorosilanes as a product is minimized. In one embodiment, the process involves contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, and a trichlorosilane. In another embodiment, the process involves contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, a trichlorosilane, and a phenylchlorosilane. In yet another embodiment, the process involves contacting a phenyl Grignard reagent, an ether solvent, an aromatic halogenated coupling solvent, and a phenylchlorosilane. In each embodiment, the reactants are present in particular mole ratios of the components.